

Anti-Protein COFACTOR ASSEMBLY OF COMPLEX C SUBUNIT B CCB2, chloroplastic antibody

Catalog: PHY2898A

Product Information

Description:	Rabbit polyclonal antibody
Background:	HCF208 is required for the biogenesis and accumulation of native cytochrome b6 in the thylakoid membrane. Controls the conversion of apocytochrome b6 to holocytochrome b6. Required for covalent binding of the c-type heme to cytochrome b6.
Synonyms:	HCF208, CCB2, COFACTOR ASSEMBLY OF COMPLEX C, HIGH CHLOROPHYLL FLUORESCENCE 208
Immunogen:	KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> HCF208 (AT5G52110).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	31 kDa
Confirmed Reactivity:	Coming soon

Research Use Only

Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Brassica napus*, *Brassica rapa*, and 80-99% homologues with the sequence in *Sorghum bicolor*, *Zea mays*, *Sorghum bicolor*, *Populus trichocarpa*. For more species homologues information, please contact tech support at tech@phytoab.com.