

Anti-Serine/threonine-protein kinase EDR1 antibody

Catalog: PHY0977S

Product Information

Description:	Rabbit polyclonal antibody
Background:	EDR1 is involved in the regulation of a MAP kinase cascade (probably including MPK3 and MPK6) that negatively regulates salicylic acid-(SA-)dependent defense responses, abscisic acid (ABA) signaling, and ethylene-induced senescence. Together with KEG, EDR1 may regulate endocytic trafficking and/or the formation of signaling complexes on trans-Golgi network (TGN)/ early endosome (EE) vesicles during stress responses.
Synonyms:	EDR1, ATEDR1, ENHANCED DISEASE RESISTANCE 1.
Immunogen:	KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> EDR1 (AT1G08720).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	104 kDa

Research Use Only

Confirmed Reactivity:

Coming soon

Predicted Reactivity:

Among species analyzed species, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica napus*, *Brassica rapa*.

For more species homologues information, please contact tech support at tech@phytoab.com.