

Anti-Histone H4 antibody

Catalog: PHY2317A

Product Information

Description: Rabbit polyclonal antibody

Background: H4 is histone 4.

Synonyms: H4, HIS4, HISTONE H4

Immunogen: KLH-conjugated synthetic peptide (16 aa from Central section) derived from

Arabidopsis thaliana H4 (AT2G28740).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 11 kDa

Confirmed Reactivity: Coming soon

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used for

immunization is 100% homologues with the sequence in *Oryza sativa*, Panicum virgatum, Zea mays, Glycine max, Brassica rapa, Hordeum vulgare, Cucumis sativus, Medicago truncatula, Vitis vinifera, Triticum

aestivum, Populus trichocarpa, Gossypium raimondii, Nicotiana


tabacum, Spinacia oleracea, Sorghum bicolor, Physcomitrium patens,



Setaria viridis.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng and 15 ng recombinant protein containing the peptide for immunization and having a molecular mass of 46 kDa.

Electrophoresis: 12% SDS-PAGE.

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.