

Anti-microtubule-associated protein 65-1/2 antibody

Catalog: PHY0917A

Product Information

Description: Rabbit polyclonal antibody

Background: The Arabidopsis thaliana MAP65-1 (AT5G55230) and MAP65-2 (AT4G26760)

genes are members of the larger eukaryotic MAP65/ASE1/PRC gene family of

microtubule-associated proteins.

Synonyms: MAP65-1, ATMAP65-1, MICROTUBULE-ASSOCIATED PROTEINS 65-1

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana MAP65-1 (AT5G55230).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 μl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 69 kDa

Confirmed Reactivity: Coming soon


Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used for

immunization is 80-99% homologues with the sequence in *Brassica* napus, *Brassica rapa*, *Gossypium raimondii*, *Solanum tuberosum*, *Populus trichocarpa*, *Spinacia oleracea*, *Panicum virgatum*, *Setaria*



viridis, Glycine max, Cucumis sativus, Nicotiana tabacum, Medicago truncatula, Solanum lycopersicum, Hordeum vulgare, Triticum aestivum. For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 5 ng and 15 ng recombinant protein containing the peptide for

immunization and having a molecular mass of 47 kDa.

Electrophoresis: 12% SDS-PAGE.

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.