

## Anti-SIr1515 protein antibody

Catalog: PHY5378S

## **Product Information**

Description:	Rabbit polyclonal antibody	
Background:	slr1515	
Synonyms:	slr1515	
Immunogen:	KLH-conjugated synthetic peptide (19 aa from C terminal section) derived from	
	Synechocystis sp. PCC 6803 slr1515.	
Form:	Lyophilized	
Quantity:	150 μg	
Purification:	Serum	
	Peptide affinity form antibody available upon request at info@phytoab.com.	
<b>Reconstitution:</b>	Reconstitution with 150 µl of sterile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ\!\!\mathbb{C}$ as supplied.	
	6 months, -20 to -70 $^\circ\!\mathbb{C}$ under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ \!$	
Shipping:	The product is shipped at 4 $^\circ\!\!{ m C}$ . Upon receipt, store it immediately at the	
	temperature recommended above.	


## **Application Information**

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	52 / 49 kDa
Confirmed Reactivity:	Synechocystis sp. PCC 6803
Predicted Reactivity:	For homologues with other species especially algae, please contact
	tech support at <u>tech@phytoab.com</u> .

**Research Use Only** 



## **Application Example**



PCC 6803-L: 30 µg whole-cell lysate protein from WT of Synechocystis *sp.* PCC 6803 (Light without Glucose).
PCC 6803-L+G: 30 µg whole-cell lysate protein from WT of Synechocystis *sp.* PCC 6803 (Light Plus Glucose).
PCC6803-D+G: 30 µg whole-cell lysate protein from WT of Synechocystis *sp.* PCC 6803 (Dark Plus Glucose).
Electrophoresis: (5-20)% Gradient gel
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4 °C.

**Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000). **Detection:** using chemiluminescence substrate and image were captured with CCD camera.

