

Anti-Alpha-amylase antibody

Catalog: PHY0004

Product Information

Description:	Rabbit polyclonal antibody
Background:	α -Amylases are hydrolytic enzymes responsible for the mobilization of the starch into metabolizable sugars. It can hydrolyze alpha bonds of large, alpha-linked polysaccharides, such as starch and glycogen, yielding glucose and maltose. This process provides the energy for the growth of roots and shoots and is crucial during germination of cereal seeds.
Synonyms:	AMY1.1, RAmy1A
Immunogen:	KLH-conjugated synthetic peptide (16 aa from N terminal section) derived from <i>Oryza sativa</i> AMY1.1 (Os02g0765600).
Form:	Lyophilized
Quantity:	150 μ g
Purification:	Protein A purified
Reconstitution:	Reconstitution with 150 μ l of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	49 kDa
Confirmed Reactivity:	<i>Hordeum vulgare</i>
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used

Research Use Only

for immunization is 100% homologues with the sequence in *Oryza sativa*, *Triticum aestivum*, *Sorghum bicolor*, *Hordeum vulgare*, and 80-99% homologues with the sequence in *Zea mays*, *Panicum virgatum*, *Setaria viridis*, *Gossypium raimondii*, *Spinacia oleracea*. For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Hv-TP: 10µg, 25 µg and 50 µg total protein from *Hordeum vulgare* leaf.

Electrophoresis: 15% SDS-Urea-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:20000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.