

## **Anti-CP47 subunit of Photosystem II antibody**

Catalog: PHY0058A

## **Product Information**

**Description:** Rabbit polyclonal antibody

Background: CP47 is one of the components of the core complex of photosystem II (PSII). It

binds chlorophyll and helps catalyze the primary light-induced photochemical

processes of PSII. CP47 is encoded by ATCG00680.

Synonyms: PsbB

**Immunogen:** KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana PsbB (ATCG00680).

Form: Lyophilized

**Quantity:** 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 56 kDa

Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in Solanum

tuberosum, Physcomitrella patens, Spinacia oleracea, Medicago


truncatula, Nicotiana tabacum, Zea mays, Populus trichocarpa, Vitis



vinifera, Brassica rapa, Setaria viridis, Gossypium raimondii, Glycine max, Cucumis sativus, Brassica napus, Sorghum bicolor, and 80-99% homologues with the sequence in Oryza sativa, Hordeum vulgare, Triticum aestivum, Leymus chinensis, Chlamydomonas reinhardtii.

For more species homologues information, please contact tech support at <a href="mailto:tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Thy-WT: thylakoid membrane protein from WT of *Arabidopsis thaliana* containing 0.25 µg of chlorophyll.

Thy-PS II mutant: thylakoid membrane protein from an Arabidopsis mutant with low accumulation of PS II containing 0.25 µg of chlorophyll.

Electrophoresis: 15% SDS- PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L (HRP) (Cat# PHY6000)

Detection: using chemiluminescence substrate and image were

captured with CCD camera.