

Anti-RNA polymerase subunit beta' antibody

Catalog: PHY1904

Product Information

Description: Mouse monoclonal (Clone: 5B1) antibody

Background: In chloroplasts, transcription of plastid genes is mediated by two types of RNA

polymerase: plastid-encoded RNA polymerase (PEP) and nuclear encoded RNA polymerase (NEP). PEP is composed of four core subunits (α , β , β ', β ") and a promoter recognition subunit (ofactor). RpoC1 (ATCG00180) is the β '

subunit of PEP.

Synonyms: RpoC1

Immunogen: Recombinant protein of RpoC1 derived from *Arabidopsis thaliana* ATCG00180.

Form: Lyophilized

Quantity: 150 μg

Purification: Protein A purified

Reconstitution: Reconstitution with 150 µl of 0.01M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected/apparent MW: 79 / 72 kDa


Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: For more species homologues information, please contact tech

support at tech@phytoab.com.



Application Example



TP: 5 µg, 10 µg and 20 µg total protein from Arabidopsis thaliana, respectively.

Electrophoresis: 12% SDS- PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:5000 dilution using Goat Anti-Rabbit IgG

H&L (HRP) (Cat# PHY6006)

Detection: using chemiluminescence substrate and image were

captured with CCD camera.