

Anti-30S ribosomal protein S1, C-terminal antibody

Catalog: PHY0424A

Product Information

Description: Rabbit polyclonal antibody

Background: Chloroplast ribosomal protein S1 (RPS1) is a heat-responsive protein and

functions in protein biosynthesis in chloroplast.

Synonyms: RPS1, ARRPS1, PLASTID RIBOSOMAL PROTEIN S1, PRPS1, RIBOSOMAL

PROTEIN S1

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana RPS1 (AT5G30510).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 45 kDa

Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in Solanum

tuberosum, Populus trichocarpa, Zea mays, Nicotiana tabacum, Brassica rapa, Solanum lycopersicum, Brassica napus, Cucumis

Research Use Only



sativus, Glycine max, Spinacia oleracea, Vitis vinifera, Setaria viridis, Oryza sativa, Sorghum bicolor, Hordeum vulgare, Panicum virgatum, Triticum aestivum, and 80-99% homologues with the sequence in Gossypium raimondii, Medicago truncatula, Physcomitrium patens, Chlamydomonas reinhardtii.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example

kDa M

Str: 15 µg stromal protein from Arabidopsis thaliana.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.