

Anti-Glucose-1-phosphate adenylyltransferase, C-terminal antibody

Catalog: PHY5159A

Product Information

Description: Rabbit polyclonal antibody

Background: GlgC is involved in the biosynthesis of ADP-glucose, a building block required

for the elongation reactions to produce glycogen. It catalyzes the reaction

between ATP and alpha-D-glucose 1-phosphate (G1P) to produce

pyrophosphate and ADP-Glc.

Synonyms: glgC, ADPGlc PPase, agp

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Synechocystis sp. PCC 6803 glgC (slr1176).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 49 kDa

Confirmed Reactivity: Synechocystis sp. PCC 6803

Predicted Reactivity: For homologues with other species especially algae, please contact



tech support at tech@phytoab.com.

Application Example

PCC 6803-TP: 15 µg and 30 µg total protein from Synechocystis sp. PCC 6803,

respectively.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:5000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.