

Anti-RNA polymerase sigma factor antibody

Catalog: PHY5207A

Product Information

Description: Rabbit polyclonal antibody

Background: Sigma factors are initiation factors that promote the attachment of RNA

polymerase to specific initiation sites and are then released.

Synonyms: sigD

Immunogen: KLH-conjugated synthetic peptide (15 aa from Central section) derived from

Synechocystis sp. PCC 6803 sigD (sll2012).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:4000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 36 kDa

Confirmed Reactivity: Synechocystis sp. PCC 6803

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 80-99% homologues with the sequence in

Synechococcus elongatus PCC 7942.

For homologues with other species especially algae, please contact



tech support at tech@phytoab.com.

Application Example

PCC 6803-TP 10μg 20μg 95 72 55 43 34 26

PHY5207A

PCC 6803-TP: 10 µg and 20 µg total protein from Synechocystis sp. PCC 6803,

respectively.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:4000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.