

## Anti-Photosystem I-associated linker protein CpcL antibody

Catalog: PHY5348S

## **Product Information**

**Description:** Rabbit polyclonal antibody

Background: cpcG2

**Synonyms:** cpcG2, Phycobilisome rod-core linker polypeptide CpcG2, cpcL

**Immunogen:** KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from

Synechocystis sp. PCC 6803 cpcG2 (sll1471).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 29 kDa

Confirmed Reactivity: Synechocystis sp. PCC 6803

Predicted Reactivity: For homologues with other species especially algae, please contact

tech support at tech@phytoab.com.



## **Application Example**

PCC 6803 kDa M L+G D+G 35— 4 PCC 6803+L/G: 30 µg whole-cell lysate protein from WT of Synechocystis

sp. PCC 6803 (Light Plus Glucose).

PCC6803+G: 30 µg whole-cell lysate protein from WT of Synechocystis

sp. PCC 6803 (Dark Plus Glucose).

Electrophoresis: (5-20)% Gradient gel

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.