

Anti-Ferredoxin C2 antibody

Catalog: PHY1147A

Product Information

Description: Rabbit polyclonal antibody

Background: FdC2 is a ferredoxin protein capable of alternative electron partitioning. FdC1

level increases in conditions of acceptor limitation at PSI.

Synonyms: FdC2, FDC2, FERREDOXIN C 2.

Immunogen: KLH-conjugated synthetic peptide (13 aa from Central section) derived from

Arabidopsis thaliana FdC2 (AT1G32550).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 22 kDa

Confirmed Reactivity: Coming soon

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in *Brassica* rapa, *Brassica napus*, and 80-99% homologues with the sequence in *Vitis vinifera*, *Gossypium raimondii*, *Populus trichocarpa*, *Medicago*

Research Use Only



truncatula, Glycine max, Solanum tuberosum, Solanum lycopersicum, Nicotiana tabacum, Panicum virgatum, Zea mays, Sorghum bicolor, Cucumis sativus, Spinacia oleracea.

For more species homologues information, please contact tech support at tech@phytoab.com.