

Anti-Arabidopsis histidine kinase 4, C-terminal antibody

Catalog: PHY0036S

Product Information

Description: Rabbit polyclonal antibody

Background: Histidine Kinases (HK) are multifunctional, typically transmembrane, proteins of

the transferase class that play a role in signal transduction across the cellular membrane. The vast majority of HKs are homodimers that exhibit autokinase, phosphotransfer, and phosphatase activity. HKs can act as cellular receptors for

signaling molecules in a way analogous to tyrosine kinase receptors (RTK).

Synonyms: AHK4, ATCRE1, CRE1, CYTOKININ RESPONSE 1, WOL, WOL1, WOODEN

LEG, WOODEN LEG 1

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana AHK4 (AT2G01830).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability & Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70℃ as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8[°]C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 118 kDa



Confirmed Reactivity: Coming soon

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 80-99% homologues with the sequence in

Brassica rapa, Brassica napus.

For more species homologues information, please contact tech

support at tech@phytoab.com.