

Anti-Phosphoglycerate kinase 1, chloroplastic antibody

Catalog: PHY0405S

Product Information

Description:	Rabbit polyclonal antibody
Background:	Phosphoglycerate kinase is an ATP-generating enzyme that acts in the glycolytic, gluconeogenic, and photosynthetic pathways. Chl-PGK is encoded in the nucleus and translated to produce a 50 kDa precursor protein and is then processed into mature 43 kDa in the chloroplast.
Synonyms:	PGK1, PHOSPHOGLYCERATE KINASE 1
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from <i>Arabidopsis thaliana</i> PGK1 (AT3G12780).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	50 / 45 kDa
Confirmed Reactivity:	<i>Arabidopsis thaliana</i>
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in <i>Brassica</i>

Research Use Only

napus, Brassica rapa, Zea mays, Triticum aestivum, Glycine max, Oryza sativa, Setaria viridis, Spinacia oleracea, Panicum virgatum, Sorghum bicolor, Medicago truncatula, Hordeum vulgare, Vitis vinifera, Populus trichocarpa, Solanum tuberosum, Gossypium raimondii, Solanum lycopersicum, Cucumis sativus.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Str: 2.75 µg, 6 µg, 12ug stromal protein from *Arabidopsis thaliana*, respectively.

Electrophoresis: 12.5% SDS-Urea-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.