

Anti-PLASTID TRANSCRIPTIONALLY ACTIVE 12 antibody

Catalog: PHY0389A

Product Information

Description:	Rabbit polyclonal antibody
Background:	In chloroplasts, transcription of plastid genes is mediated by two types of RNA polymerase: plastid-encoded RNA polymerase (PEP) and nuclearencoded RNA polymerase (NEP). Transcription in plastids is also mediated by a number of nuclear-encoded factors in addition to PEP and NEP. In the insoluble RNA polymerase preparation samples, a total of 18 components named as pTACs (pTAC1 to pTAC18) were identified. pTAC12 (AT2G34640) is one of the components associated with PEP complex.
Synonyms:	pTAC12, HEMERA, HMR, PLASTID TRANSCRIPTIONALLY ACTIVE 12, TAC12
Immunogen:	KLH-conjugated synthetic peptide (15 aa from Central section) derived from <i>Arabidopsis thaliana</i> pTAC12 (AT2G34640).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
------------------------------	--

Research Use Only

Expected/apparent MW: 61 kDa

Confirmed Reactivity: Coming soon

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Solanum tuberosum*, *Brassica napus*, *Medicago truncatula*, *Nicotiana tabacum*, *Solanum lycopersicum*, *Gossypium raimondii*, *Brassica rapa*, and 80-99% homologues with the sequence in *Vitis vinifera*, *Cucumis sativus*, *Glycine max*, *Spinacia oleracea*, *Populus trichocarpa*, *Oryza sativa*, *Triticum aestivum*, *Hordeum vulgare*, *Zea mays*, *Setaria viridis*, *Panicum virgatum*.
For more species homologues information, please contact tech support at tech@phytoab.com.