

# Anti-RNA polymerase subunit beta antibody

Catalog: PHY1239

### **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** In chloroplasts, transcription of plastid genes is mediated by two types of RNA

polymerase: plastid-encoded RNA polymerase (PEP) and nuclear encoded

RNA polymerase (NEP). PEP is composed of four core subunits ( $\alpha$ ,  $\beta$ , $\beta$ ', $\beta$ ") and

a promoter recognition subunit (ofactor). RpoB (ATCG00190) is the  $\boldsymbol{\beta}$  subunit of

PEP.

Synonyms: RpoB, RNA POLYMERASE SUBUNIT BETA

Immunogen: Recombinant protein of RpoB (537-1072 aa) derived from Arabidopsis thaliana

ATCG00190.

Form: Lyophilized

**Quantity:** 150 μg

Purification: Protein A

**Reconstitution:** Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

### **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 121 kDa


Confirmed Reactivity: Arabidopsis thaliana

**Predicted Reactivity:** For more species homologues information, please contact tech



### support at tech@phytoab.com.

## **Application Example**



TP: 5 µg, 10 µg, and 20 µg total protein from Arabidopsis thaliana, respectively.

Electrophoresis: 10% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at  $4^{\circ}$ C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.