

Anti-Abscisic acid 8'-hydroxylase 2, C-terminal antibody

Catalog: PHY1832S

Product Information

Description: Rabbit polyclonal antibody

Background: CYP707A2 is a protein with ABA 8'-hydroxylase activity, involved in ABA

catabolism. It appears to play a major role in the rapid decrease in ABA levels

during early seed imbibition.

Synonyms: CYP707A2, "CYTOCHROME P450, FAMILY 707, SUBFAMILY A,

POLYPEPTIDE 2"

Immunogen: KLH-conjugated synthetic peptide (18 aa from C terminal section) derived from

Arabidopsis thaliana CYP707A2 (AT2G29090).

Form: Lyophilized

Quantity: 150 μg

Purification: Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 55 kDa

Confirmed Reactivity: Coming soon



Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Brassica rapa, Brassica napus*, and 80-99% homologues with the sequence in *Gossypium raimondii, Populus trichocarpa, Spinacia oleracea, Glycine max, Cucumis sativus.*

For more species homologues information, please contact tech support at tech@phytoab.com.