

Anti-PsaE subunit of photosystem I antibody

Catalog: PHY3112A

Product Information

Description: Rabbit polyclonal antibody

Background: PsaE is part of the stromal side of photosystem I (PSI). In *Arabidopsis*

thaliana, the two nuclear genes PsaE1 and PsaE2 code for PSI-E, and

transcripts of PsaE1 are markedly more abundant than PsaE2 transcripts.

Synonyms: PsaE

Immunogen: KLH-conjugated synthetic peptide (14 aa from Central section) derived from

Arabidopsis thaliana PsaE (AT4G28750).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 15 / 14 kDa

Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide

used for immunization is 100% homologues with the sequence in

Medicago truncatula, Brassica rapa, Brassica napus, and 80-99%


homologues with the sequence in Glycine max, Populus trichocarpa,



Gossypium raimondii, Vitis vinifera, Solanum tuberosum, Solanum lycopersicum, Spinacia oleracea, Cucumis sativus.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Thy-WT: thylakoid membrane protein from WT of *Arabidopsis thaliana* containing 2.5 µg of chlorophyll.

Thy-PS II: thylakoid membrane protein from an Arabidopsis mutant with low accumulation of PS II containing 2.5 µg of chlorophyll.

Electrophoresis: Tricine-SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.