

Anti-Lhca4 protein of LHCI antibody

Catalog: PHY0084A

Product Information

Description: Rabbit polyclonal antibody

Background: The light-harvesting complex (LHC) functions as a light receptor, it captures

and delivers excitation energy to photosystem with which it is closely

associated. The light-harvesting protein Lhca4 is one of the four main and

highly conserved types of chlorophyll a/b-binding proteins (Lhca1-4) of the light

harvesting antenna (LHCI) of plant photosystem I. Lhca4 is imported as a

precursor from the cytosol into the chloroplast. Upon insertion into the thylakoid

membrane Lhca4 forms a heterodimer (LHCI-730) with Lhca1 that associates

with the PSI core close to PsaG and PsaF.

Synonyms: Lhca4, CAB4, LHCA4, LIGHT-HARVESTING CHLOROPHYLL-PROTEIN

COMPLEX I SUBUNIT A4

Immunogen: KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from

Arabidopsis thaliana Lhca4 (AT3G47470).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability & Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.



Expected / apparent MW: 28 / 23 kDa

Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in

Medicago truncatula.

For more species homologues information, please contact tech

support at tech@phytoab.com.

Application Example



Thy: Thylakoid membrane protein from Arabidopsis thaliana containing 0.1 µg and

0.25 µg of chlorophyll, respectively.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitroceOllulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000)

Detection: using chemiluminescence substrate and image were captured with

CCD camera.