

Anti-BIOTIN/LIPOYL ATTACHMENT DOMAIN CONTAINING 2 antibody

Catalog: PHY1765S

Product Information

Description: Rabbit polyclonal antibody

Background: BADC2 is a single hybrid motif superfamily protein. BADC2 significantly inhibits

the activity of Acetyl-CoA carboxylase (ACCase) in Arabidopsis thaliana.

Synonyms: BADC2, BCCP-LIKE PROTEIN 1, BIOTIN/LIPOYL ATTACHMENT DOMAIN

CONTAINING 2, BLP1

Immunogen: KLH-conjugated synthetic peptide (14 aa from N terminal section) derived from

Arabidopsis thaliana BADC2 (AT1G52670).

Form: Lyophilized

Quantity: 150 μg

Purification: Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 30 / 24 kDa


Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: For more species homologues information, please contact tech



support at tech@phytoab.com.

Application Example



Str: 6 µg and 12µg stromal protein from Arabidopsis thaliana, respectively.

Electrophoresis: 15% SDS-Urea-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.