

## Anti-Chloroplastic group IIA intron splicing facilitator CRS1, chloroplastic antibody

Catalog: PHY6310S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** CRS1 is required for the splicing of group IIA introns in chloroplasts, and

especially for atpF, by regulating the intron folding. And it is involved in

chloroplast protein translation.

**Synonyms**: CRS1

Immunogen: Recombinant protein of CRS1 derived from Zea mays CRS1

(Zm00001d032373).

Form: Lyophilized

Quantity:150 μgPurification:Serum

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 81 / 72 kDa


Confirmed Reactivity: Zea mays

**Predicted Reactivity:** For more species homologues information, please contact tech

support at tech@phytoab.com.



## **Application Example**



Zm-TP: 15 µg and 30 µg total protein from Zea mays.

Electrophoresis: 10% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.