

Anti-Heat shock protein antibody

Catalog: PHY5324S

Product Information

Description:	Rabbit polyclonal antibody					
Background:	htpG					
Synonyms:	htpG					
Immunogen:	KLH-conjugated synthetic peptide (16 aa from Central section) derived from					
	Synechocystis sp. PCC 6803 htpG (sll0430).					
Form:	Lyophilized					
Quantity:	150 μg					
Purification:	Serum					
	Peptide affinity form antibody available upon request at info@phytoab.com.					
Reconstitution:	ution: Reconstitution with 150 µl of sterile water.					
	"Note: please spin tube briefly prior to opening it to avoid any losses that might					
	occur from lyophilized material adhering to the cap or sides of the tube".					
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.					
Storage:	12 months from date of receipt, -20 to -70°C as supplied.					
	6 months, -20 to -70°C under sterile conditions after reconstitution.					
	1 month, 2 to 8°C under sterile conditions after reconstitution.					
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the					
	temperature recommended above.					

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)				
	Note: Optimal dilutions/concentrations should be determined by the				
	end user.				
Expected / apparent MW:	75 / 83 kDa				
Confirmed Reactivity:	Synechocystis sp. PCC 6803				
Predicted Reactivity:	For homologues with other species especially algae, please contact				
	tech support at tech@phytoab.com.				

Research Use Only



Application Example

		PCC 6803-TP				
kDa	M	L	L+G	D+G		
98				-		
					<	
75						
62						
48				-		
32						
		DUN	153240	2		

L: 30 µg whole-cell lysate protein from WT of *Synechocystis sp.* PCC 6803 (Light without Glucose). L+G: 30 µg whole-cell lysate protein from WT of *Synechocystis sp.* PCC 6803 (Light Plus Glucose). G: 30 µg whole-cell lysate protein from WT of *Synechocystis sp.* PCC 6803 (Dark Plus Glucose). Electrophoresis: (5-20)% Gradient gel Transfer: blotting to NC (nitrocellulose) membrane for 1 h. Blocking: 5% skim milk at RT or 4°C for 1 h. Primary antibody: 1:1000 dilution overnight at 4°C. Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000). Detection: using chemiluminescence substrate and image were captured with CCD camera.

PHY5324S

Research Use Only