

Anti-Hypersensitive to High Light 1, C-terminal antibody

Catalog: PHY0483S

Product Information

Description:	Rabbit polyclonal antibody
Background:	HHL1 is a chloroplast protein and participates in photodamage repair of PSII under high light. Mutants without HHL1 mutants show hypersensitivity to high light and drastically decreased PSII photosynthetic activity. HHL1 associates with the PSII core monomer complex through direct interaction with PSII core proteins CP43 and CP47, and also interacts with LQY1, which functions in the repair and reassembly of PSII.
Synonyms:	HHL1, HYPERSENSITIVE TO HIGH LIGHT 1
Immunogen:	KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> HHL1 (AT1G67700).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	26 / 24 kDa

Research Use Only

Confirmed Reactivity: *Arabidopsis thaliana*

Predicted Reactivity: For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Thy-WT: thylakoid membrane protein from *Arabidopsis thaliana* containing 2.5 µg, and 5 µg of chlorophyll, respectively.

Thy-*hhl1*: thylakoid membrane protein from a *hhl1* mutant of *Arabidopsis thaliana* containing 5 µg of chlorophyll.

Electrophoresis: 15% SDS-Urea-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.