

Anti-Histone H3 antibody

Catalog: PHY2460A

Product Information

Description: Rabbit polyclonal antibody

Background: Histone H3 is one of the five main histone proteins involved in the structure of

chromatin in eukaryotic cells. H3 is involved with the structure of the

nucleosomes of the 'beads on a string' structure. Histone H3 is an important protein in the emerging field of epigenetics, where its sequence variants and variable modification states are thought to play a role in the dynamic and long

term regulation of genes.

Synonyms: H3

Immunogen: KLH-conjugated synthetic peptide (14 aa from N terminal section) derived from

Arabidopsis thaliana H3.3 (AT4G40030, AT4G40040, AT5G10980) and Histone

H3-like 3 (AT1G75600).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 15 kDa



Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in *Glycine* max, *Hordeum vulgare*, *Solanum tuberosum*, *Zea mays*, *Spinacia* oleracea, *Triticum aestivum*, *Sorghum bicolor*, *Panicum virgatum*,

Physcomitrella patens, Chlamydomonas reinhardtii, Cucumis sativus,

Oryza sativa, Medicago truncatula, Gossypium raimondii, Vitis vinifera, Setaria viridis, Brassica rapa, Populus trichocarpa, and 80-99% homologues with the sequence in Nicotiana tabacum,

Brassica napus.

The sequence of the synthetic peptide used for immunization is 93% homologues with the sequence in Histone H3-like 1 (AT1G13370), Histone H3.2 (AT5G65360, AT5G10400, AT5G10390, AT3G27360, AT1G09200), Histone H3-like 2 (AT1G19890), and Histone H3-like 4 (AT5G12910).

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example

kDa M

10

PHY2460A

Nuc

Nuc: 1.5 µg nuclear protein from *Arabidopsis thaliana*.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.