

Anti-3-dehydroquinate synthase antibody

Catalog: PHY2780A

Product Information

Description:	Rabbit polyclonal antibody
Background:	AT3G28760
Synonyms:	AT3G28760
Immunogen:	KLH-conjugated synthetic peptide of AT3G28760 (16aa from central section) derived from <i>Arabidopsis thaliana</i> (AT3G28760).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	47 / 44 kDa
Confirmed Reactivity:	<i>Arabidopsis thaliana</i>
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in <i>Brassica rapa</i> , <i>Glycine max</i> , <i>Brassica napus</i> , <i>Cucumis sativus</i> , <i>Nicotiana tabacum</i> , <i>Solanum tuberosum</i> , <i>Oryza sativa</i> , <i>Solanum lycopersicum</i> , and 80-99% homologues with the sequence in <i>Vitis vinifera</i> ,

Research Use Only

Medicago truncatula, *Zea mays*, *Hordeum vulgare*, *Sorghum bicolor*,
Setaria viridis, *Panicum virgatum*, *Triticum aestivum*, *Populus*
trichocarpa, *Spinacia oleracea*, *Chlamydomonas reinhardtii*,
Physcomitrium patens, *Gossypium raimondii*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Chl: 7.5 µg total chloroplast protein from *Arabidopsis thaliana*.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.