

## Anti-HIGH CHLOROPHYLL FLUORESCENCE 106 antibody

Catalog: PHY2202S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** HCF106 is the thylakoid membrane delta pH translocation pathway component

protein, it is related to Escherichia coli TatA and TatB. It is involved in delta

pH-dependent protein transport required for chloroplast development,

especially thylakoid membrane formation.

Synonyms: HCF106, HIGH CHLOROPHYLL FLUORESCENCE 106, TWIN-ARGININE

TRANSLOCATION B, Sec-independent protein translocase protein TATB,

chloroplastic

**Immunogen:** Recombinant protein of HCF106 (107-260 aa) derived from *Arabidopsis* 

thaliana AT5G52440.

Form: Lyophilized

Quantity:150 μgPurification:Serum

**Reconstitution:** Reconstitution with 150µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 28 kDa



Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: For more species homologues information, please contact tech

support at tech@phytoab.com.

## **Application Example**

Thy: thylakoid membrane protein from  $Arabidopsis\ thaliana$  containing 5  $\mu g$  of chlorophyll.

Electrophoresis: 15% SDS-Urea-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000)

**Detection:** using chemiluminescence substrate and image were captured with CCD

camera.