

Anti-Chaperonin 60 subunit alpha 2, chloroplastic antibody

Catalog: PHY1358S

Product Information

Description:	Rabbit polyclonal antibody
Background:	Type I chaperonins are large, double ring complexes involved in mediating the folding of unfolded proteins. In chloroplasts, chaperonin complex is designated Cpn60 complex. In Arabidopsis, two Cpn60 α subunits and four Cpn60 β subunits are present. Cpn60 α 2 (AT5G18820) is one of putative Cpn60 α subunit with low expression level in chloroplasts and is essential for embryo development.
Synonyms:	CPN60A2, CHAPERONIN-60ALPHA2, CPNA2, CPN60ALPHA2, EMB3007, EMBRYO DEFECTIVE 3007
Immunogen:	KLH-conjugated synthetic peptide (16 aa from N terminal section) derived from <i>Arabidopsis thaliana</i> CPN60A2 (AT5G18820).
Form:	Lyophilized
Quantity:	150 μ g
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 μ l of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
------------------------------	--------------------------------------------------------------------------------------------------------------

Research Use Only

Expected / apparent MW: 61 kDa

Confirmed Reactivity: *Arabidopsis thaliana*

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica rapa*, *Brassica napus*.
For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Chl: 7.5 µg total chloroplast protein from *Arabidopsis thaliana*.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP) (Cat# PHY6000)

Detection: using chemiluminescence substrate and image were captured with CCD camera.