

Anti-Cyclin-dependent kinase F-1, C-terminal antibody

Catalog: PHY1421S

Product Information

Description: Rabbit polyclonal antibody

Background: Plants have two distinct types of CAKs, namely CDKD and CDKF; in

Arabidopsis, CDKF;1 exhibits the highest CDK kinase activity in vitro. CDKF;1

also functions in the activation of CDKD;2 and CDKD;3 by T-loop

phosphorylation.

Synonyms: Cak1At, CAK1AT, CDK-ACTIVATING KINASE 1AT, CDKF;1,

CYCLIN-DEPENDENT KINASE F;1

Immunogen: KLH-conjugated synthetic peptide (11 aa from C terminal section) derived from

Arabidopsis thaliana Cak1At (AT4G28980).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70 $^{\circ}$ C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 54 kDa

Confirmed Reactivity: Coming soon

Research Use Only



Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Brassica* rapa, *Brassica* napus.

For more species homologues information, please contact tech support at tech@phytoab.com.