

Anti-Peptidyl-prolyl cis-trans isomerase FKBP16-4, chloroplastic, C-terminal antibody

Catalog: PHY2773A

Product Information

Description: Rabbit polyclonal antibody

Background: AT3G10060

Synonyms: AT3G10060, Peptidyl-prolyl cis-trans isomerase FKBP16-4, chloroplastic,

PPlase FKBP16-4, FK506-binding protein 16-4, AtFKBP16-4, Immunophilin

FKBP16-4, Rotamase, FKBP16-4, FKBP24-2

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana AT3G10060.

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 25 / 16 kDa

Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in Brassica



napus, Brassica rapa, Triticum aestivum, Panicum virgatum, Oryza sativa, Gossypium raimondii, Nicotiana tabacum, Solanum tuberosum, Solanum lycopersicum, Sorghum bicolor, Hordeum vulgare, Zea mays, Setaria viridis, and 80-99% homologues with the sequence in Physcomitrium patens, Vitis vinifera, Glycine max, Cucumis sativus, Spinacia oleracea, Populus trichocarpa, Medicago truncatula.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Information



Chl: 7.5 µg total chloroplast protein from Arabidopsis thaliana.

TP: 30 µg total protein from Arabidopsis thaliana.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 2 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgGH&L(HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image werecaptured with

CCD camera.