

Anti-Glyceraldehyde-3-phosphate dehydrogenase GAPC, cytosolic antibody

Catalog: PHY3070S

Product Information

Description:	Rabbit polyclonal antibody
Background:	Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) is an enzyme of ~37kDa that catalyzes the sixth step of glycolysis and thus serves to break down glucose for energy and carbon molecules. Plants contain both cytosolic and chloroplastic GAPDHs (glyceraldehyde-3-phosphate dehydrogenases). In <i>Arabidopsis thaliana</i> , cytosolic GAPDH is involved in the glycolytic pathway and is represented by two differentially expressed isoforms (GAPC1 AT3G04120 and GAPC2 AT1G13440) that are 98% identical in amino acid sequence.
Synonyms:	GAPDH, GAPC, GAPC, GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE C
Immunogen:	KLH-conjugated synthetic peptide (17 aa from N terminal section) derived from <i>Arabidopsis thaliana</i> GAPC1(AT3G04120) and GAPC2 (AT1G13440).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:4000) Note: Optimal dilutions/concentrations should be determined by the
------------------------------	--

Research Use Only

Expected / apparent MW:

end user.
37 kDa


Confirmed Reactivity:

Arabidopsis thaliana

Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Solanum tuberosum*, *Glycine max*, *Brassica napus*, *Brassica rapa*, *Medicago truncatula*, *Oryza sativa*, *Vitis vinifera*, *Spinacia oleracea*, *Zea mays*, *Triticum aestivum*, *Gossypium raimondii*, *Physcomitrium patens*, *Populus trichocarpa*, *Nicotiana tabacum*, *Solanum lycopersicum*.
For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Cyto: 15 µg and 30 µg cytosolic protein from *Arabidopsis thaliana*.
TP: 15 µg and 30 µg total protein from *Arabidopsis thaliana*.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:4000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG &L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.