

Anti-Uroporphyrinogen decarboxylase 2, chloroplastic antibody

Catalog: PHY2680S

Product Information

Description:	Rabbit polyclonal antibody
Background:	HEME2 catalyzes the decarboxylation of four acetate groups of uroporphyrinogen-III to yield coproporphyrinogen-III.
Synonyms:	HEME2
Immunogen:	KLH-conjugated synthetic peptide (21 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> HEME2 (AT2G40490).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:5000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	44 / 40 kDa
Confirmed Reactivity:	<i>Arabidopsis thaliana</i>
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in <i>Hordeum</i>

Research Use Only

vulgare, *Populus trichocarpa*, *Glycine max*, *Nicotiana tabacum*, *Spinacia oleracea*, *Vitis vinifera*, *Cucumis sativus*, *Solanum tuberosum*, *Brassica rapa*, *Oryza sativa*, *Brassica napus*, *Solanum lycopersicum*, *Gossypium raimondii*, and 80-99% homologues with the sequence in *Zea mays*, *Sorghum bicolor*, *Panicum virgatum*, *Physcomitrium patens*, *Setaria viridis*, *Medicago truncatula*.
For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Chl: 5 µl total chloroplast protein from *Arabidopsis thaliana*.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:5000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000)

Detection: using chemiluminescence substrate and image were captured with CCD camera.