

Anti-Thioredoxin-like fold domain-containing protein MRL7L, chloroplastic antibody

Catalog: PHY2505A

Product Information

Description: Rabbit polyclonal antibody

Background: MRL7-L plays an essential role in early steps of chloroplast development. It's

required for the proper function of the plastid transcriptional machinery and

protein accumulation in thylakoid membranes.

Synonyms: MRL7-L, MESOPHYLL-CELL RNAI LIBRARY LINE 7-LIKE

Immunogen: KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from

Arabidopsis thaliana MRL7-L (AT2G31840).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 41 / 30 kDa

Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in Brassica




rapa, Brassica napus, and 80-99% homologues with the sequence in Nicotiana tabacum, Solanum tuberosum, Solanum lycopersicum, Populus trichocarpa, Glycine max, Vitis vinifera.

For more species homologues information, please contact tech

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Chl: 7.5 µg total chloroplast protein from Arabidopsis thaliana.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.