

## Anti-NdhV, chloroplastic antibody

Catalog: PHY0307

## **Product Information**

**Description:** Rabbit polyclonal antibody

Background: Chloroplast NADH dehydrogenase-like (NDH) complex mediates cyclic

electron transport around PSI and chlororespiration. Chloroplast NDH consists of five subcomplexes. In higher plants, NDH complex further associates with photosystem I (PSI) to form the NDH-PSI supercomplex. NdhV is a subunit of

the NDH complex.

Synonyms: NdhV

Immunogen: Recombinant mature NdhV Protein without chloroplast targeting peptide

derived from Arabidopsis thaliana AT2G04039.

Form: Lyophilized

**Quantity:** 150 μg

Purification: Protein A purified

**Reconstitution:** Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 21 / 18 kDa


Confirmed Reactivity: Arabidopsis thaliana

**Predicted Reactivity:** For more species homologues information, please contact tech

support at tech@phytoab.com.



## **Application Example**



Chl: 7.5 µg total chloroplast protein from Arabidopsis thaliana.

Thy: thylakoid membrane protein from *Arabidopsis thaliana* containing 2.5 µg of chlorophyll.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with

CCD camera.