

Anti-PLASTID TRANSCRIPTIONALLY ACTIVE 2 antibody

Catalog: PHY2528A

Product Information

Description:	Rabbit polyclonal antibody
Background:	In chloroplasts, transcription of plastid genes is mediated by two types of RNA polymerase: plastid-encoded RNA polymerase (PEP) and nuclear encoded RNA polymerase (NEP). Transcription in plastids is also mediated by a number of nuclear-encoded factors in addition to PEP and NEP. In the insoluble RNA polymerase preparation samples, a total of 18 components named as pTACs (pTAC1 to pTAC18) were identified. pTAC2 (AT1G74850) is one of the components associated with PEP complex.
Synonyms:	pTAC2, PDE343, PIGMENT DEFECTIVE 343, PLASTID TRANSCRIPTIONALLY ACTIVE 2, PTAC2
Immunogen:	KLH-conjugated synthetic peptide (16 aa from N terminal section) derived from <i>Arabidopsis thaliana</i> pTAC2 (AT1G74850).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
------------------------------	--

Research Use Only

Expected / apparent MW:

96 / 75 kDa

Confirmed Reactivity:


Arabidopsis thaliana, *Oryza sativa*, *Zea mays*

Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Brassica napus*, *Glycine max*, *Vitis vinifera*, *Sorghum bicolor*, *Zea mays*, *Setaria viridis*, *Panicum virgatum*, *Oryza sativa*, *Brassica rapa*, *Cucumis sativus*, *Medicago truncatula*, *Solanum lycopersicum*, *Nicotiana tabacum*, *Solanum tuberosum*, *Gossypium raimondii*, and 80-99% homologues with the sequence in *Hordeum vulgare*, *Triticum aestivum*, *Spinacia oleracea*, *Physcomitrium patens*, *Populus trichocarpa*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



At-Chl: 80 µg and 10 µl total chloroplast protein from *Arabidopsis thaliana*, respectively.

Zm-Chl: 5 µl total chloroplast protein from *Zea mays*.

Os-Chl: 5 µl total chloroplast protein from *Oryza sativa*.

Electrophoresis: 10% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000)

Detection: using chemiluminescence substrate and image were captured with CCD camera.