

Anti-Lhca6 protein of LHCI antibody

Catalog: PHY0470S

Product Information

| Description: | Rabbit polyclonal antibody | |
|------------------------|---|--|
| Background: | The light-harvesting complex (LHC) functions as a light receptor, it captures | |
| | and delivers excitation energy to photosystems with which it is closely | |
| | associated. Lhca6 seems involved in the function of the photosystem I in low | |
| | light conditions, when other LHCA proteins are less abundant. Required, | |
| | together with LHCA5, for the formation of a full-size NAD(P)H | |
| | dehydrogenase-photosystem I supercomplex (NDH-PSI) that triggers cyclic | |
| | and chlororespiratory electron transport in chloroplast thylakoids, especially | |
| | under stress conditions. | |
| Synonyms: | Lhca6, LHCA6, LHCA2*1, PHOTOSYSTEM I LIGHT HARVESTING | |
| | COMPLEX GENE 6 | |
| Immunogen: | KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from | |
| | Arabidopsis thaliana Lhca6 (AT1G19150). | |
| Form: | Lyophilized | |
| Quantity: | 150 µg | |
| Purification: | Serum | |
| | Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> . | |
| Reconstitution: | Reconstitution with 150 µl of sterile water. | |
| | "Note: please spin tube briefly prior to opening it to avoid any losses that might | |
| | occur from lyophilized material adhering to the cap or sides of the tube". | |
| Stability & | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. | |
| Storage: | 12 months from date of receipt, -20 to -70 $^\circ C$ as supplied. | |
| | 6 months, -20 to -70 $^\circ C$ under sterile conditions after reconstitution. | |
| | 1 month, 2 to 8 $^\circ\!{ m C}$ under sterile conditions after reconstitution. | |
| Shipping: | The product is shipped at $4^\circ\!\mathbb{C}$. Upon receipt, store it immediately at the | |
| | temperature recommended above. | |
| | | |

Application Information

| Recommended Dilution: | Western Blot (1:1000-1:2000) |
|------------------------------|--|
| | Note: Optimal dilutions/concentrations should be determined by the |

Research Use Only



end user.

Expected / apparent MW: Confirmed Reactivity: Predicted Reactivity: 30 kDa

Coming soon

Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica napus, Brassica rapa, Glycine max.* For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.



Research Use Only