Anti-PENETRATION 2, C-terminal antibody

Catalog: PHY1785S

Product Information

Description:	Rabbit polyclonal antibody
Background:	PEN2 is a glycosyl hydrolase that localizes to peroxisomes and acts as a component of an inducible preinvasion resistance mechanism. PEN2 is a tail-anchored protein with dual-membrane targeting to peroxisomes and mitochondria and that PEN2 has the capacity to form homo-oligomer complexes.
Synonyms:	PEN2, BETA GLUCOSIDASE 26, BGLU26, PENETRATION 2
Immunogen:	KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from *Arabidopsis thaliana* PEN2 (AT2G44490).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com.
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70℃ as supplied. 6 months, -20 to -70℃ under sterile conditions after reconstitution. 1 month, 2 to 8℃ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4℃. Upon receipt, store it immediately at the temperature recommended above.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	64 / 58 kDa
Confirmed Reactivity:	*Arabidopsis thaliana*

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Brassica rapa, Brassica napus.*

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



TP: 20 μg, 40 μg and 80 μg total protein from *Arabidopsis thaliana*, respectively.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP) (Cat# PHY6000)

Detection: using chemiluminescence substrate and image were captured with CCD camera.