

Anti-Translocase of chloroplast 159, chloroplastic antibody

Catalog: PHY0467A

Product Information

Description: Rabbit polyclonal antibody

Background: TOC159 is an integral membrane GTPase that functions as a transit-sequence

receptor required for the import of proteins necessary for chloroplast

biogenesis. It located in the outer chloroplast membrane.

Synonyms: TOC159, ATTOC159, PLASTID PROTEIN IMPORT 2, PPI2, TOC160, TOC86,

TRANSLOCON AT THE OUTER ENVELOPE MEMBRANE OF

CHLOROPLASTS 159, TRANSLOCON AT THE OUTER ENVELOPE

MEMBRANE OF CHLOROPLASTS 160, TRANSLOCON AT THE OUTER

ENVELOPE MEMBRANE OF CHLOROPLASTS 86

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana TOC159 (AT4G02510).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 161 kDa




Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica rapa*, *Brassica napus*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 5 ng ,15 ng and 50 ng recombinant protein containing the peptide for immunization and having a molecular mass of 55 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.