

# Anti-Protein TIC56, chloroplastic antibody

Catalog: PHY1632S

### **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** TIC56 is involved in protein precursor import into chloroplasts. It may be part of

an intermediate translocation complex acting as a protein-conducting channel

at the inner envelope.

Synonyms: TIC56, TRANSLOCON AT THE INNER ENVELOPE MEMBRANE OF

**CHLOROPLASTS 56** 

Immunogen: KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from

Arabidopsis thaliana TIC56 (AT5G01590).

Form: Lyophilized

Quantity:150 μgPurification:Serum

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 62 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 80-99% homologues with the sequence in


Brassica rapa, Brassica napus.

For more species homologues information, please contact tech



#### support at tech@phytoab.com.

### **Application Example**



Recom: 2.5 ng and 10 ng recombinant protein containing the peptide for

immunization and having a molecular mass of 70 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000)

**Detection:** using chemiluminescence substrate and image were captured with

CCD camera.