

Anti-NADH dehydrogenase subunit 7antibody

Catalog: PHY1077A

Product Information

Description:	Rabbit polyclonal antibody	
Background:	Complex I is the largest protein complex of the oxidative phosphorylation	
	system in mitochondrial and it catalyzes NADH-quinone oxidoreduction.	
	Complex I represents the main entrance site for electrons into the respiratory	
	electron transfer chain. In Arabidopsis, Complex I have at least 49 subunits	
	and NAD7 (ATMG00510) is one of the subunit.	
Synonyms:	NAD7, NADH DEHYDROGENASE SUBUNIT 7	
Immunogen:	unogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from	
	Arabidopsis thaliana NAD7 (ATMG00510).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Immunogen affinity purified	
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.	
	6 months, -20 to -70 $^\circ C$ under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ$ C under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	45 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide
	used for immunization is 100% homologues with the sequence in
	Vitis vinifera, Brassica napus, Setaria viridis, Triticum aestivum,

PhytoAB Inc.



Glycine max, Medicago truncatula, Oryza sativa, Gossypium raimondii, Zea mays, rassica rapa, Hordeum vulgare. For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were

captured with CCD camera.

PHY1077A

Research Use Only