

Anti-Ferrochelatase-1 antibody

Catalog: PHY2901S

Product Information

Description: Rabbit polyclonal antibody

Background: Ferrochelatase I catalyzes the last step of heme biosynthesis by inserting

ferrous iron into protoporphyrin IX to produce protoheme.

Synonyms: FC1, ATFC-I, FC-I, FERROCHELATASE 1, FUS3-COMPLEMENTING GENE 1

Immunogen: KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from

Arabidopsis thaliana FC1 (AT5G26030).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 52 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in *Brassica* napus, *Brassica* rapa, and 80-99% homologues with the sequence in *Sorghum bicolor*, *Zea mays*, *Cucumis sativus*, *Nicotiana tabacum*, *Glycine max*, *Triticum aestivum*, *Populus trichocarpa*, *Solanum*


Research Use Only



tuberosum, Solanum lycopersicum, Oryza sativa, Medicago truncatula, Hordeum vulgare, Spinacia oleracea, Setaria viridis, Panicum virgatum, Vitis vinifera, Gossypium raimondii.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 15 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.