

Anti-Protein COFACTOR ASSEMBLY OF COMPLEX C SUBUNIT B CCB3, chloroplastic antibody

Catalog: PHY2899S

Product Information

Description: Rabbit polyclonal antibody

Background: CCB3 is one of four Arabidopsis homologs of bacterial ymlg proteins. It is

required for the biogenesis and accumulation of native cytochrome b6 in the

thylakoid membrane. It controls the conversion of apocytochrome b6 to

holocytochrome b6. It is required for covalent binding of the c-type heme to

cytochrome b6.

Synonyms: CCB3, ATYLMG3, COFACTOR ASSEMBLY, COMPLEX C (B6F), YLMG3

Immunogen: KLH-conjugated synthetic peptide (18 aa from N terminal section) derived from

Arabidopsis thaliana CCB3 (AT5G36120).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 19 kDa




Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Brassica rapa*, *Brassica napus*, and 80-99% homologues with the sequence in *Spinacia oleracea*, *Gossypium raimondii*, *Glycine max*, *Medicago truncatula*, *Cucumis sativus*, *Populus trichocarpa*, *Vitis vinifera*, *Solanum tuberosum*, *Solanum lycopersicum*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 15 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.