

Anti-Chlorophyll a-b binding protein 2, chloroplastic, N-terminal antibody

Catalog: PHY3287S

Product Information

Description:	Rabbit polyclonal antibody
Background:	The light-harvesting chlorophyll a/b-binding proteins of photosystem II (LHCII)
	are the major components of the photosynthetic machinery in plants which
	contain more than 60% of plant chlorophyll.
	The LHCII proteins can be grouped into six subfamilies (Lhcb1-6) which are
	encoded by LHC gene family, Lhcb1, Lhcb2 and Lhcb3 are the major
	pigment-binding proteins which are encoded by Lhcb1, Lhcb2 and Lhcb3
	genes, respectively. Lhcb1, Lhcb2 and Lhcb3 polypeptides each with about 232
	amino acid residues are similar in sequence, Lhcb1-3 precursors are
	synthesized in cytoplasm and following transport into chloroplasts inserted into
	thylakoid membranes structure and function. Lhcb1 and Lhcb2 are the most
	abundant proteins in the light harvesting antenna complex.
Synonyms:	Lhcb2, Lhcb2.1/2.2/2.3
Immunogen:	KLH-conjugated synthetic peptide (14 aa from N terminal section) derived from
	Arabidopsis thaliana Lhcb2.1 (AT2G05100), Lhcb2.2 (AT2G05070), and
	Lhcb2.3 (AT3G27690).
Form:	Lyophilized
Quantity:	150 μg
Purification:	Serum
	Peptide affinity form antibody available upon request at info@phytoab.com.
Reconstitution:	Reconstitution with 150 µl of sterile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.
	6 months, -20 to -70 $^\circ\!\!\!\mathrm{C}$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the
	temperature recommended above.

Research ese enty




Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	29 / 25 kDa
Confirmed Reactivity:	Arabidopsis thaliana
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in <i>Brassica</i>
	napus, Brassica rapa, Vitis vinifera, and 80-99% homologues with the
	sequence in Hordeum vulgare subsp. vulgare, Solanum tuberosum,
	Solanum lycopersicum, Cucumis sativus, Nicotiana tabacum, Glycine
	max, Populus trichocarpa, Oryza sativa Japonica Group, Panicum
	virgatum, Sorghum bicolor, Setaria viridis, Triticum aestivum,
	Medicago truncatula, Zea mays, Spinacia oleracea, Physcomitrium
	patens, Gossypium raimondii .
	For more species homologues information, please contact tech
	support at <u>tech@phytoab.com</u> .

Application Example

Example1




Thy: Thylakoid membrane protein from *Arabidopsis thaliana* leaf containing 0.1 μg and 0.25μg of chlorophyll, respectively. **Electrophoresis:** 15% SDS-PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:2000 dilution overnight at 4°C. **Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000). **Detection:** using chemiluminescence substrate and image were captured with

CCD camera.

Research Use Only



Example2



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 48 kDa. **Electrophoresis:** 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only