

Anti-Malate dehydrogenase [NADP], chloroplastic, C-terminal antibody

Catalog: PHY3968S

Product Information

Description: Rabbit polyclonal antibody

Background: NADP-MDH belongs to lactate/malate dehydrogenase family protein;

Synonyms: NADP-MDH, NADP-DEPENDENT MALATE DEHYDROGENASE

Immunogen: KLH-conjugated synthetic peptide (17 aa from C terminal section) derived from

Arabidopsis thaliana NADP-MDH (AT5G58330).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4 °C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 48 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in Brassica

napus, Glycine max, Brassica rapa, Vitis vinifera, Populus

trichocarpa, Gossypium raimondii, Medicago truncatula, Spinacia



oleracea, Hordeum vulgare, Oryza sativa, Triticum aestivum, Solanum lycopersicum, and 80-99% homologues with the sequence in Solanum tuberosum, Nicotiana tabacum, Cucumis sativus, Setaria viridis, Panicum virgatum, Sorghum bicolor, Zea mays.

For more species homologues information, please contact tech support at tech@phytoab.com.