

Anti-PsbW subunit of Photosystem II, C-terminal antibody

Catalog: PHY3519A

Product Information

Description: Rabbit polyclonal antibody

Background: PsbW is a nuclear-encoded PSII subunit and found in higher plants but not in

cyanobacteria. PsbW has an apparent molecular mass of 6.1 kDa containing

54 amino acids.

Synonyms: PsbW, Photosystem II reaction center W protein, chloroplastic, PSII 6.1 kDa

protein

Immunogen: KLH-conjugated synthetic peptide (12 aa from C terminal section) derived from

Arabidopsis thaliana PsbW (AT2G30570).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4 °C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 14 / 9 kDa


Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used



for immunization is 100% homologues with the sequence in *Glycine max, Gossypium raimondii,* and 80-99% homologues with the sequence in *Cucumis sativus, Brassica rapa, Brassica napus, Nicotiana tabacum, Vitis vinifera, Populus trichocarpa, Medicago truncatula, Spinacia oleracea, Sorghum bicolor, Oryza sativa.*For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



WT: thylakoid membrane protein from WT of *Arabidopsis thaliana* containing 2.5 µg of chlorophyll.

PSII mutant: thylakoid membrane protein from an Arabidopsis mutant with low accumulation of PSI containing 2.5 µg of chlorophyll.

Electrophoresis: Tricine-SDS-PAGE.

Transfer: blotting to NC (nitrocellulose) membrane for 1h.

Blocking: 5% skim milk at RT or 4°C for 1h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.