

Anti-NdhO, chloroplastic antibody

Catalog: PHY3268S

Product Information

Description:	Rabbit polyclonal antibody		
-			
Background:	Chloroplast NADH dehydrogenase-like (NDH) complex mediates cyclic		
	electron transport around PSI and chlororespiration. Chloroplast NDH consists		
	of five subcomplexes. In higher plants, NDH complex further associates with		
	photosystem I (PSI) to form the NDH-PSI supercomplex. NdhO is a subunit of		
	the NDH complex.		
Synonyms:	NdhO, NAD(P)H dehydrogenase subunit O, NADH-plastoquinone oxidoreductase subunit O, NDH subunit O		
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from <i>Arabidopsis thaliana</i> NdhO (AT1G74880).		
Form:	Lyophilized		
Quantity:	150 µg		
Purification:	Serum		
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .		
Reconstitution:	Reconstitution with 150 µl of sterile water.		
	"Note: please spin tube briefly prior to opening it to avoid any losses that migh		
	occur from lyophilized material adhering to the cap or sides of the tube".		
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.		
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb C$ as supplied.		
	6 months, -20 to -70 $^\circ\!{ m C}$ under sterile conditions after reconstitution.		
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.		
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the		
	temperature recommended above.		

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)	
	Note: Optimal dilutions/concentrations should be determined by	/ the
	end user.	
Expected / apparent MW:	18 kDa	



Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica napus*, *Brassica rapa*.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 48 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only