

Anti-Nitrile-specifier protein 5 antibody

Catalog: PHY2516S

Product Information

Description: Rabbit polyclonal antibody

Background: NSP5 is a nitrile-specifier protein. NSP5 is one out of five (At3g16400/NSP1,

At2g33070/NSP2, At3g16390/NSP3, At3g16410/NSP4 and At5g48180/NSP5)

A. thaliana epithiospecifier protein (ESP) homologues that promote simple

nitrile, but not epithionitrile or thiocyanate formation.

Synonyms: NSP5, ATNSP5, NITRILE SPECIFIER PROTEIN 5

Immunogen: KLH-conjugated synthetic peptide (14 aa from N terminal section) derived from

Arabidopsis thaliana NSP5 (AT5G48180).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4° C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 36 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in

Brassica napus, Brassica rapa.



For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 40 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with

CCD camera.