

Anti-Aspartic proteinase A1 antibody

Catalog: PHY1683S

Product Information

Description: Rabbit polyclonal antibody

Background: APA1 is an aspartic proteinase that forms a heterodimer and is stable over a

broad pH range (ph 3-8).

Synonyms: APA1, ASPARTIC PROTEINASE A1, ATAPA1, ATPASPA1, PASPA1,

PUTATIVE ASPARTIC PROTEINASE A1

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana APA1 (AT1G11910).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 55 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in *Brassica* napus, *Brassica* rapa, *Glycine* max, and 80-99% homologues with the sequence in *Triticum* aestivum, *Hordeum* vulgare, *Populus*



trichocarpa, Zea mays, Oryza sativa, Spinacia oleracea, Panicum virgatum, Medicago truncatula, Setaria viridis.

The sequence of the synthetic peptide used for immunization is 93% (14/15) homologues with the sequence in PASPA2 (AT1G62290). For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 48 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with

CCD camera.