

Anti-C-phycocyanin beta chain antibody

Catalog: PHY5243A

Product Information

Description: Rabbit polyclonal antibody

Background: Phycocyanin beta subunit, encoded by the CpcB (sl/1577) in Synechocystis sp.

PCC 6803.

Synonyms: CpcB

Immunogen: KLH-conjugated synthetic peptide (14 aa from N terminal section) derived from

Synechocystis sp. PCC 6803 CpcB (sll1577).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 $^{\circ}$ C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 18 kDa

Confirmed Reactivity: Synechocystis sp. PCC 6803

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 80-99% homologues with the sequence in


Synechococcus elongatus PCC 7942.

For more species homologues information, please contact tech



support at tech@phytoab.com.

Application Example



PCC 6803-TP: 31.25 μg and 62.5 μg total protein from Synechocystis sp. PCC

6803, respectively.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.