

## Anti-Plasmodesmata-located protein 5 antibody

Catalog: PHY1993A

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** PDLP5 is a plasmodesmal protein that may be involved in the intercellular

movement of molecules through the plasmodesmata. The protein has two

DUF26 domains and a single transmembrane domain.

**Synonyms:** PDLP5, HOPW1-1-INDUCED GENE1, HWI1, PLASMODESMATA-LOCATED

**PROTEIN 5** 

Immunogen: KLH-conjugated synthetic peptide (15 aa from Central section) derived from

Arabidopsis thaliana PDLP5 (AT1G70690).

Form: Lyophilized

**Quantity:** 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 μl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at  $4^{\circ}$ C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.


Expected / apparent MW: 32 kDa

**Predicted Reactivity:** For more species homologues information, please contact tech

support at tech@phytoab.com.



## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 40 kDa.

inimunization and having a molecular mass of 40 kDa

**Electrophoresis:** 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4℃.

**Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with

CCD camera.